ELSEVIER

Brain Research 755 (1997) 356–358

BRAIN RESEARCH

Author Index

Abbadie, C., see Peterson, M.A. (755) 9
Abdul-Ghani, A.-S., Attwell, P.J.E., Singh Kent, N., Bradford, H.F., Croucher, M.J. and Jane, D.E.
Anti-epileptogenic and anticonvulsant activity of L-2-amino-4-phosphonobutyrate, a presynaptic glutamate receptor agonist (755) 202
Abrahamsen, G.C. and Carr, K.D.
Effect of adrenalectomy on cocaine facilitation of lateral hypothalamic self-stimulation (755) 156
Aghajanian, G.K., see Jolas, T. (755) 229
Ahmed, B.Y., see Tokuda, M. (755) 162
Arai, T., see Chang, M.C.J. (755) 74
Attwell, P.J.E., see Abdul-Ghani, A.-S. (755) 202

Baik, H.H., see Cho, S. (755) 335
Barnes, J.R., see Raley-Susman, K.M. (755) 1
Basbaum, A.I., see Peterson, M.A. (755) 9
Bell, J.M., see Chang, M.C.J. (755) 74
Benes, F.M., Wickramasinghe, R., Vincent, S.L., Khan, Y. and Todtenkopf, M.
Uncoupling of $GABA_A$ and benzodiazepine receptor binding activity in the hippocampal formation of schizophrenic brain (755) 121
Bittencourt, J.C., see Elias, C.F. (755) 255
Blanchard, J.H., see Meyer-Bernstein, E.L. (755) 112
Bradford, H.F., see Abdul-Ghani, A.-S. (755) 202
Bredow, S., see Obál Jr., F. (755) 130
Bremer, E.G., see Yamamoto, H. (755) 175
Buijs, R.M., see Van der Beek, E.M. (755) 101

Carr, K.D., see Abrahamsen, G.C. (755) 156
Carter, J.G., see McDougal Jr., D.B. (755) 304
Cazevieille, C. and Osborne, N.N.
Retinal neurones containing kainate receptors are influenced by exogenous kainate and ischaemia while neurones lacking these receptors are not — melatonin counteracts the effect of ischaemia and kainate (755) 91
Cerullo, L., see Yamamoto, H. (755) 175
Chad, J.E., see Pringle, A.K. (755) 36
Chang, M.C.J., Arai, T., Freed, L.M., Wakabayashi, S., Channing, M.A., Dunn, B.B., Der, M.G., Bell, J.M., Sasaki, T., Herscovitch, P., Eckelman, W.C. and Rapoport, S.I.
Brain incorporation of [1-^{11}C]arachidonate in normocapnic and hypercapnic monkeys, measured with positron emission tomography (755) 74
Channing, M.A., see Chang, M.C.J. (755) 74
Chatterjee, S.S., see Klein, J. (755) 347
Chen, J., Koyama, N., Kaneko, T. and Mizuno, N.
The locus coeruleus of the Japanese monkey (*Macaca fuscata*) does not express μ-opioid receptor-like immunoreactivity (755) 326
Chi, M.M.-Y., see McDougal Jr., D.B. (755) 304
Cho, S., Joh, T.H., Baik, H.H., Dibinis, C. and Volpe, B.T.
Melatonin administration protects CA1 hippocampal neurons after transient forebrain ischemia in rats (755) 335
Christensen, B.N., see Giblin, L.J. (755) 146
Colling, S.B., Khana, M., Collinge, J. and Jefferys, J.G.R.
Mossy fibre reorganization in the hippocampus of prion protein null mice (755) 28
Collinge, J., see Colling, S.B. (755) 28
Copray, J.C.V.M., see Liem, R.S.B. (755) 319
Cowsette, B.R., see McDougal Jr., D.B. (755) 304
Croucher, M.J., see Abdul-Ghani, A.-S. (755) 202

Dean, R.G., see Edwards, G.L. (755) 84
Der, M.G., see Chang, M.C.J. (755) 74
Dibinis, C., see Cho, S. (755) 335
Dirnagl, U., see Hock, C. (755) 293
Dunn, B.B., see Chang, M.C.J. (755) 74
Dworkin, S.I., see Martin, T.J. (755) 313

Eckelman, W.C., see Chang, M.C.J. (755) 74
Edwards, G.L., White, B.D., He, B., Dean, R.G. and Martin, R.J.
Elevated hypothalamic neuropeptide Y levels in rats with dorsomedial hindbrain lesions (755) 84
Elias, C.F. and Bittencourt, J.C.
Study of the origins of melanin-concentrating hormone and neuropeptide EI immunoreactive projections to the periaqueductal gray matter (755) 255
Elliott, E.J., Parks, D.A. and Fishman, P.S.
Effect of proximal axotomy on GAP-43 expression in cortical neurons in the mouse (755) 221

Farabaugh, S.M., see Wild, J.M. (755) 137
Fdez. Espejo, E. and Gil, E.
Single restraint stress sensitizes acute chewing movements induced by haloperidol, but not if the 5-HT_{1A} agonist 8-OH-DPAT is given prior to stress (755) 351
Feuer, P., Loeb, A., Monitto, C.L., O'Rourke, M. and Kurth, C.D.
Cerebral blood flow regulation in neonatal rabbits is altered by chronic cocaine administration (755) 142
Fishman, P.S., see Elliott, E.J. (755) 221
Fox, G.Q. and Kriebel, M.E.
Dynamic responses of presynaptic terminal membrane pools following KCl and sucrose stimulation (755) 47
Freed, L.M., see Chang, M.C.J. (755) 74
Furuse, M., Matsumoto, M., Okumura, J.-i., Sugahara, K. and Hasegawa, S.
Intracerebroventricular injection of mammalian and chicken glucagon-like peptide-1 inhibits food intake of the neonatal chick (755) 167

Giblin, L.J. and Christensen, B.N.
Connexin43 immunoreactivity in the catfish retina (755) 146
Gil, E., see Fdez. Espejo, E. (755) 351
Guha-Thakurta, N., see Obál Jr., F. (755) 130

Hama, A.T., see Siegan, J.B. (755) 331
Hasbroucq, T., see Romaiguère, P. (755) 181
Hasegawa, S., see Furuse, M. (755) 167
Hashimoto, Y., Hori, H., Kawakami, T., Kusakabe, T. and Takenaka, T.
Effect of dibutyryl cyclic AMP on axoplasmic transport in the hippocampus (755) 343
Hatase, O., see Tokuda, M. (755) 162
He, B., see Edwards, G.L. (755) 84
Heekeren, H., see Hock, C. (755) 293
Herscovitch, P., see Chang, M.C.J. (755) 74
Hock, C., Villringer, K., Müller-Spahn, F., Wenzel, R., Heekeren, H., Schuh-Hofer, S., Hofmann, M., Minoshima, S., Schwaiger, M., Dirnagl, U. and Villringer, A.
Decrease in parietal cerebral hemoglobin oxygenation during performance of a verbal fluency task in patients with Alzheimer's disease monitored by means of near-infrared spectroscopy (NIRS) — correlation with simultaneous rCBF-PET measurements (755) 293
Hofmann, M., see Hock, C. (755) 293
Hori, H., see Hashimoto, Y. (755) 343
Htain, W.-W., Leong, S.-K., Yuen, R. and Ling, E.-A.
Effects of thymosin α1 on the development of amoeboid microglial cells in the corpus callosum of neonatal BALB/c and athymic mice (755) 63

ELSEVIER

Brain Research 755 (1997) 356–358

BRAIN RESEARCH

Author Index

Abbadie, C., see Peterson, M.A. (755) 9
Abdul-Ghani, A.-S., Attwell, P.J.E., Singh Kent, N., Bradford, H.F., Croucher, M.J. and Jane, D.E.
Anti-epileptogenic and anticonvulsant activity of L-2-amino-4-phosphonobutyrate, a presynaptic glutamate receptor agonist (755) 202
Abrahamsen, G.C. and Carr, K.D.
Effect of adrenalectomy on cocaine facilitation of lateral hypothalamic self-stimulation (755) 156
Aghajanian, G.K., see Jolas, T. (755) 229
Ahmed, B.Y., see Tokuda, M. (755) 162
Arai, T., see Chang, M.C.J. (755) 74
Attwell, P.J.E., see Abdul-Ghani, A.-S. (755) 202

Baik, H.H., see Cho, S. (755) 335
Barnes, J.R., see Raley-Susman, K.M. (755) 1
Basbaum, A.I., see Peterson, M.A. (755) 9
Bell, J.M., see Chang, M.C.J. (755) 74
Benes, F.M., Wickramasinghe, R., Vincent, S.L., Khan, Y. and Todtenkopf, M.
Uncoupling of $GABA_A$ and benzodiazepine receptor binding activity in the hippocampal formation of schizophrenic brain (755) 121
Bittencourt, J.C., see Elias, C.F. (755) 255
Blanchard, J.H., see Meyer-Bernstein, E.L. (755) 112
Bradford, H.F., see Abdul-Ghani, A.-S. (755) 202
Bredow, S., see Obál Jr., F. (755) 130
Bremer, E.G., see Yamamoto, H. (755) 175
Buijs, R.M., see Van der Beek, E.M. (755) 101

Carr, K.D., see Abrahamsen, G.C. (755) 156
Carter, J.G., see McDougal Jr., D.B. (755) 304
Cazevieille, C. and Osborne, N.N.
Retinal neurones containing kainate receptors are influenced by exogenous kainate and ischaemia while neurones lacking these receptors are not — melatonin counteracts the effect of ischaemia and kainate (755) 91
Cerullo, L., see Yamamoto, H. (755) 175
Chad, J.E., see Pringle, A.K. (755) 36
Chang, M.C.J., Arai, T., Freed, L.M., Wakabayashi, S., Channing, M.A., Dunn, B.B., Der, M.G., Bell, J.M., Sasaki, T., Herscovitch, P., Eckelman, W.C. and Rapoport, S.I.
Brain incorporation of [1-^{11}C]arachidonate in normocapnic and hypercapnic monkeys, measured with positron emission tomography (755) 74
Channing, M.A., see Chang, M.C.J. (755) 74
Chatterjee, S.S., see Klein, J. (755) 347
Chen, J., Koyama, N., Kaneko, T. and Mizuno, N.
The locus coeruleus of the Japanese monkey (*Macaca fuscata*) does not express μ-opioid receptor-like immunoreactivity (755) 326
Chi, M.M.-Y., see McDougal Jr., D.B. (755) 304
Cho, S., Joh, T.H., Baik, H.H., Dibinis, C. and Volpe, B.T.
Melatonin administration protects CA1 hippocampal neurons after transient forebrain ischemia in rats (755) 335
Christensen, B.N., see Giblin, L.J. (755) 146
Colling, S.B., Khana, M., Collinge, J. and Jefferys, J.G.R.
Mossy fibre reorganization in the hippocampus of prion protein null mice (755) 28
Collinge, J., see Colling, S.B. (755) 28
Copray, J.C.V.M., see Liem, R.S.B. (755) 319
Cowsette, B.R., see McDougal Jr., D.B. (755) 304
Croucher, M.J., see Abdul-Ghani, A.-S. (755) 202

Dean, R.G., see Edwards, G.L. (755) 84
Der, M.G., see Chang, M.C.J. (755) 74
Dibinis, C., see Cho, S. (755) 335
Dirnagl, U., see Hock, C. (755) 293
Dunn, B.B., see Chang, M.C.J. (755) 74
Dworkin, S.I., see Martin, T.J. (755) 313

Eckelman, W.C., see Chang, M.C.J. (755) 74
Edwards, G.L., White, B.D., He, B., Dean, R.G. and Martin, R.J.
Elevated hypothalamic neuropeptide Y levels in rats with dorsomedial hindbrain lesions (755) 84
Elias, C.F. and Bittencourt, J.C.
Study of the origins of melanin-concentrating hormone and neuropeptide EI immunoreactive projections to the periaqueductal gray matter (755) 255
Elliott, E.J., Parks, D.A. and Fishman, P.S.
Effect of proximal axotomy on GAP-43 expression in cortical neurons in the mouse (755) 221

Farabaugh, S.M., see Wild, J.M. (755) 137
Fdez. Espejo, E. and Gil, E.
Single restraint stress sensitizes acute chewing movements induced by haloperidol, but not if the 5-HT_{1A} agonist 8-OH-DPAT is given prior to stress (755) 351
Feuer, P., Loeb, A., Monitto, C.L., O'Rourke, M. and Kurth, C.D.
Cerebral blood flow regulation in neonatal rabbits is altered by chronic cocaine administration (755) 142
Fishman, P.S., see Elliott, E.J. (755) 221
Fox, G.Q. and Kriebel, M.E.
Dynamic responses of presynaptic terminal membrane pools following KCl and sucrose stimulation (755) 47
Freed, L.M., see Chang, M.C.J. (755) 74
Furuse, M., Matsumoto, M., Okumura, J.-i., Sugahara, K. and Hasegawa, S.
Intracerebroventricular injection of mammalian and chicken glucagon-like peptide-1 inhibits food intake of the neonatal chick (755) 167

Giblin, L.J. and Christensen, B.N.
Connexin43 immunoreactivity in the catfish retina (755) 146
Gil, E., see Fdez. Espejo, E. (755) 351
Guha-Thakurta, N., see Obál Jr., F. (755) 130

Hama, A.T., see Siegan, J.B. (755) 331
Hasbroucq, T., see Romaiguère, P. (755) 181
Hasegawa, S., see Furuse, M. (755) 167
Hashimoto, Y., Hori, H., Kawakami, T., Kusakabe, T. and Takenaka, T.
Effect of dibutyryl cyclic AMP on axoplasmic transport in the hippocampus (755) 343
Hatase, O., see Tokuda, M. (755) 162
He, B., see Edwards, G.L. (755) 84
Heekeren, H., see Hock, C. (755) 293
Herscovitch, P., see Chang, M.C.J. (755) 74
Hock, C., Villringer, K., Müller-Spahn, F., Wenzel, R., Heekeren, H., Schuh-Hofer, S., Hofmann, M., Minoshima, S., Schwaiger, M., Dirnagl, U. and Villringer, A.
Decrease in parietal cerebral hemoglobin oxygenation during performance of a verbal fluency task in patients with Alzheimer's disease monitored by means of near-infrared spectroscopy (NIRS) — correlation with simultaneous rCBF-PET measurements (755) 293
Hofmann, M., see Hock, C. (755) 293
Hori, H., see Hashimoto, Y. (755) 343
Htain, W.-W., Leong, S.-K., Yuen, R. and Ling, E.-A.
Effects of thymosin α1 on the development of amoeboid microglial cells in the corpus callosum of neonatal BALB/c and athymic mice (755) 63

Iannotti, F., see Pringle, A.K. (755) 36
Iijima, M., see Sasaki, A. (755) 193
Inglis, W.L. and Semba, K.
Discriminable excitotoxic effects of ibotenic acid, AMPA, NMDA and quinolinic acid in the rat laterodorsal tegmental nucleus (755) 17
Isobe, Y. and Nishino, H.
GABAergic control of Arg-vasopressin release from suprachiasmatic nucleus slice culture (755) 213

Jane, D.E., see Abdul-Ghani, A.-S. (755) 202
Jefferys, J.G.R., see Colling, S.B. (755) 28
Jiang, Z.D., Moore, D.R. and King, A.J.
Sources of subcortical projections to the superior colliculus in the ferret (755) 279
Joh, T.H., see Cho, S. (755) 335
Jolas, T. and Aghajanian, G.K.
Opioids suppress spontaneous and NMDA-induced inhibitory postsynaptic currents in the dorsal raphe nucleus of the rat in vitro (755) 229

Kacsóh, B., see Obál Jr., F. (755) 130
Kaja, J., see Raley-Susman, K.M. (755) 1
Kalra, S.P., see Sahu, A. (755) 339
Kaneko, T., see Chen, J. (755) 326
Kaneko, Y., see Yamamoto, H. (755) 175
Kawakami, T., see Hashimoto, Y. (755) 343
Kersey, D., see Yamamoto, H. (755) 175
Khan, Y., see Benes, F.M. (755) 121
Khana, M., see Colling, S.B. (755) 28
Kim, A.J., see Shiozaki, T. (755) 170
King, A.J., see Jiang, Z.D. (755) 279
Kishi, T., see Tsumori, T. (755) 151
Klein, J., Chatterjee, S.S. and Löffelholz, K.
Phospholipid breakdown and choline release under hypoxic conditions: inhibition by bilobalide, a constituent of *Ginkgo biloba* (755) 347
Konishi, R., see Tokuda, M. (755) 162
Koyama, N., see Chen, J. (755) 326
Kriebel, M.E., see Fox, G.Q. (755) 47
Krueger, J.M., see Obál Jr., F. (755) 130
Kurth, C.D., see Feuer, P. (755) 142
Kusakabe, T., see Hashimoto, Y. (755) 343

Laitinen, J.T., see Laitinen, K.S.M. (755) 272
Laitinen, K.S.M., Salovaara, K., Severgnini, S. and Laitinen, J.T.
Regulation of cyclic GMP levels in the rat frontal cortex in vivo: effects of exogenous carbon monoxide and phosphodiesterase inhibition (755) 272
Leestma, J., see Yamamoto, H. (755) 175
Leong, S.-K., see Htain, W.-W. (755) 63
Liem, R.S.B., Copray, J.C.V.M. and Van der Want, J.J.L.
Dopamine-immunoreactivity in the rat mesencephalic trigeminal nucleus: an ultrastructural analysis (755) 319
Ling, E.-A., see Htain, W.-W. (755) 63
Loeb, A., see Feuer, P. (755) 142
Löffelholz, K., see Klein, J. (755) 347
Lowry, O.H., see McDougal Jr., D.B. (755) 304
Lu, Y.-F., see Tokuda, M. (755) 162

Manchester, J.K., see McDougal Jr., D.B. (755) 304
Martin, R.J., see Edwards, G.L. (755) 84
Martin, T.J., Miller Jr., M., Dworkin, S.I., Smith, J.E. and Porrino, L.J.
Alteration of local cerebral glucose utilization following intravenous administration of heroin in Fischer 344 rats (755) 313
Matsui, H., see Tokuda, M. (755) 162
Matsumoto, M., see Furuse, M. (755) 167
McDougal Jr., D.B., Cowsette, B.R., Pusateri, M.E., Carter, J.G., Manchester, J.K., Chi, M.M.-Y. and Lowry, O.H.
Glutamate and potassium stimulation of hippocampal slices metabolizing glucose or glucose and pyruvate (755) 304
McKay, W.R., see Peterson, M.A. (755) 9
Meyer-Bernstein, E.L., Blanchard, J.H. and Morin, L.P.
The serotonergic projection from the median raphe nucleus to the suprachiasmatic nucleus modulates activity phase onset, but not other circadian rhythm parameters (755) 112
Miller Jr., M., see Martin, T.J. (755) 313
Minoshima, S., see Hock, C. (755) 293
Miyamoto, O., see Tokuda, M. (755) 162
Mizuno, N., see Chen, J. (755) 326
Mkrdichian, E., see Yamamoto, H. (755) 175
Monitto, C.L., see Feuer, P. (755) 142
Moore, D.R., see Jiang, Z.D. (755) 279
Morin, L.P., see Meyer-Bernstein, E.L. (755) 112
Moskal, J.R., see Yamamoto, H. (755) 175
Müller-Spahn, F., see Hock, C. (755) 293

Nakazato, Y., see Sasaki, A. (755) 193
Nishino, H., see Isobe, Y. (755) 213

Obál Jr., F., Kacsóh, B., Bredow, S., Guha-Thakurta, N. and Krueger, J.M.
Sleep in rats rendered chronically hyperprolactinemic with anterior pituitary grafts (755) 130
Okumura, J.-i., see Furuse, M. (755) 167
Ono, K., see Tsumori, T. (755) 151
O'Rourke, M., see Feuer, P. (755) 142
Osborne, N.N., see Cazevieille, C. (755) 91

Parks, D.A., see Elliott, E.J. (755) 221
Peterson, M.A., Basbaum, A.I., Abbadie, C., Rohde, D.S., McKay, W.R. and Taylor, B.K.
The differential contribution of capsaicin-sensitive afferents to behavioral and cardiovascular measures of brief and persistent nociception and to Fos expression in the formalin test (755) 9
Porrino, L.J., see Martin, T.J. (755) 313
Possamaï, C.-A., see Romaiguère, P. (755) 181
Pringle, A.K., Iannotti, F., Wilde, G.J.C., Chad, J.E., Seeley, P.J. and Sundstrom, L.E.
Neuroprotection by both NMDA and non-NMDA receptor antagonists in in vitro ischemia (755) 36
Pusateri, M.E., see McDougal Jr., D.B. (755) 304

Raley-Susman, K.M., Barnes, J.R. and Kaja, J.
Effects of volatile anesthetics on hippocampal slice metabolism, response to anoxia with and without glucose (755) 1
Rapoport, S.I., see Chang, M.C.J. (755) 74
Reinke, H., see Wild, J.M. (755) 137
Rohde, D.S., see Peterson, M.A. (755) 9
Romaiguère, P., Possamaï, C.-A. and Hasbroucq, T.
Motor cortex involvement during choice reaction time: a transcranial magnetic stimulation study in man (755) 181
Rusak, B., see Ying, S.-W. (755) 246

Sagen, J., see Siegan, J.B. (755) 331
Sahu, A., Sninsky, C.A. and Kalra, S.P.
Evidence that hypothalamic neuropeptide Y gene expression and NPY levels in the paraventricular nucleus increase before the onset of hyperphagia in experimental diabetes (755) 339
Saito, T., see Yamamoto, H. (755) 175
Salovaara, K., see Laitinen, K.S.M. (755) 272
Sasaki, A., Iijima, M., Yokoo, H., Shoji, M. and Nakazato, Y.
Human choroid plexus is an uniquely involved area of the brain in amyloidosis: a histochemical, immunohistochemical and ultrastructural study (755) 193
Sasaki, T., see Chang, M.C.J. (755) 74
Schuh-Hofer, S., see Hock, C. (755) 293
Schwaiger, M., see Hock, C. (755) 293
Seeley, P.J., see Pringle, A.K. (755) 36
Semba, K., see Inglis, W.L. (755) 17
Severgnini, S., see Laitinen, K.S.M. (755) 272
Shiozaki, T., Taneda, M., Yasuda, M. and Kim, A.J.
Asymmetrical regulation of blood pressure in the rat medulla (755) 170
Shoji, M., see Sasaki, A. (755) 193
Siegan, J.B., Hama, A.T. and Sagen, J.
Suppression of neuropathic pain by a naturally-derived peptide with NMDA antagonist activity (755) 331
Singh Kent, N., see Abdul-Ghani, A.-S. (755) 202
Smith, J.E., see Martin, T.J. (755) 313
Sninsky, C.A., see Sahu, A. (755) 339
Sugahara, K., see Furuse, M. (755) 167
Sundstrom, L.E., see Pringle, A.K. (755) 36

Takenaka, T., see Hashimoto, Y. (755) 343
Taneda, M., see Shiozaki, T. (755) 170
Taylor, B.K., see Peterson, M.A. (755) 9
Todtenkopf, M., see Benes, F.M. (755) 121
Tokuda, M., Ahmed, B.Y., Lu, Y.-F., Matsui, H., Miyamoto, O., Yamaguchi, F., Konishi, R. and Hatase, O.
Involvement of calmodulin-dependent protein kinases-I and -IV in long-term potentiation (755) 162
Tsumori, T., Ono, K., Kishi, T. and Yasui, Y.
Demonstration of the corticotectobulbar pathway from the orofacial motor cortex to

the parvicellular reticular formation in the rat (755) 151

Van den Hurk, R., see Van der Beek, E.M. (755) 101
Van der Beek, E.M., Wiegant, V.M., Van Oudheusden, H.J.C., Van der Donk, H.A., Van den Hurk, R. and Buijs, R.M. Synaptic contacts between gonadotropin-releasing hormone-containing fibers and neurons in the suprachiasmatic nucleus and perichiasmatic area: an anatomical substrate for feedback regulation? (755) 101
Van der Donk, H.A., see Van der Beek, E.M. (755) 101
Van der Want, J.J.L., see Liem, R.S.B. (755) 319
Van Oudheusden, H.J.C., see Van der Beek, E.M. (755) 101
Villringer, A., see Hock, C. (755) 293
Villringer, K., see Hock, C. (755) 293
Vincent, S.L., see Benes, F.M. (755) 121
Volpe, B.T., see Cho, S. (755) 335

Wakabayashi, S., see Chang, M.C.J. (755) 74
Wenzel, R., see Hock, C. (755) 293
White, B.D., see Edwards, G.L. (755) 84
Wickramasinghe, R., see Benes, F.M. (755) 121
Wiegant, V.M., see Van der Beek, E.M. (755) 101
Wild, J.M., Reinke, H. and Farabaugh, S.M. A non-thalamic pathway contributes to a whole body map in the brain of the budgerigar (755) 137
Wilde, G.J.C., see Pringle, A.K. (755) 36

Yamaguchi, F., see Tokuda, M. (755) 162
Yamamoto, H., Saito, T., Kaneko, Y., Kersey, D., Yong, V.W., Bremer, E.G., Mkrdichian, E., Cerullo, L., Leestma, J. and Moskal, J.R. α2,3-Sialyltransferase mRNA and α2,3-linked glycoprotein sialylation are increased in malignant gliomas (755) 175
Yasuda, M., see Shiozaki, T. (755) 170
Yasui, Y., see Tsumori, T. (755) 151
Ying, S.-W. and Rusak, B. 5-HT_7 receptors mediate serotonergic effects on light-sensitive suprachiasmatic nucleus neurons (755) 246
Yokoo, H., see Sasaki, A. (755) 193
Yong, V.W., see Yamamoto, H. (755) 175
Yuen, R., see Htain, W.-W. (755) 63